

Travelling Waves in Hamiltonian Systems on 2D Lattices with Nearest Neighbor Interactions*

Michal Fečkan[†] and Vassilis M. Rothos^{††}

[†] Department of Mathematical Analysis and Numerical Mathematics, Comenius University,
Mlynská dolina, 842 48 Bratislava, Slovakia,
and Mathematical Institute, Slovak Academy of Sciences,
Štefánikova 49, 814 73 Bratislava, Slovakia

^{††} Department of Mathematics, Physics and Computational Sciences,
Faculty of Technology, Aristotle University of Thessaloniki, Thessaloniki 54124, Greece

February 8, 2008

Abstract

We study travelling waves on a two-dimensional lattice with linear and nonlinear coupling between nearest particles and a periodic nonlinear substrate potential. Such a discrete system can model molecules adsorbed on a substrate crystal surface. We show the existence of both uniform sliding states and periodic travelling waves as well in a two-dimensional sine-Gordon lattice equation using topological and variational methods.

1 Introduction

Nonlinear lattice models are used in many physical applications [35]. Spatial discreteness is obviously the standard situation for applications in solid state physics. One of the well-known class of discrete models with a broad variety of applications, consists of the sine-Gordon dynamical lattices, also known as Frenkel–Kontorova (FK) models, e.g. [30]. For instance, the FK model is a model for the motion of a dislocation in a crystal, for an adsorbate layer on the surface of a crystal, for ionic conductors, for glassy materials, for charge-density-wave transport, for chains of coupled Josephson junctions, and for sliding friction as well [38]. Another important discrete models are either discrete nonlinear Schrödinger systems [35] used in a variety of applications ranging from nonlinear optics to the theory of molecular vibrations, or a discrete model of Fermi, Pasta and Ulam (FPU) [19, 20] as a prototypical case of nonlinear mass-spring chains. Systems of discrete lattices are also studied. For instance, two coupled one-dimensional

*Nonlinearity 2007(accepted)

(1d) nonlinear lattices are used in [31] to represent the Bernal-Fowler filaments in ice or more complex biological macromolecules in membranes.

Most lattice systems are not integrable. On the other hand, there is an important class of completely integrable lattice equations, such as the Toda lattice, the Ablowitz–Ladik lattice, and the integrable discrete sine-Gordon lattice [12, 35]. These lattices possess exact soliton solutions showing that discreteness of the host media does not preclude the propagation of localized coherent structures. Discrete solitons and more specifically kink-like topological excitations are ubiquitous structures that arise in numerous physical applications ranging from dislocations or ferroelectric domain walls in solids, to bubbles in DNA, or magnetic chains and Josephson junctions, among others [1, 16]. Hence one of the most important problems in discrete systems is the existence of travelling waves and uniform sliding states [37, 38]. Of course, the study of their stability is also crucial.

Recently, particular interest has been devoted to the dynamics of structures on two-dimensional (2d) lattice systems. For instance [11, 13], coherent localized and extended defects such as dislocations, domains walls, vortices, grain boundaries, etc., play an important role in the dynamical properties of materials with applications to the problem of adsorbates deposited on crystal surfaces, in superlattices of ultrathin layers, or in large-area Josephson junctions, which are described as 2d FK type models. On the other hand, a 2d cubic Hamiltonian lattice appears in elastostatic investigation [18] modeling a particles interaction via interatomic potentials, which is a natural 1d analogy of the 1d FPU lattice.

In this paper, we focus on undamped 2d FK models and their generalizations. Motivated by [39], we consider an isotropic two-dimensional planar model where rigid molecules rotate in the plane of a square lattice. At site (n, m) the angle of rotation is $u_{n,m}$, each molecule interacts linearly with its first nearest neighbors and with a nonlinear periodic substrate potential. Here, G is the linear coupling coefficient and ω_0^2 is the strength of the potential barrier or square of the frequency of small oscillations in the bottom of the potential wells. The equation of motion of the rotator at site (n, m) is

$$\ddot{u}_{n,m} = G[u_{n+1,m} + u_{n-1,m} - 2u_{n,m} + u_{n,m+1} + u_{n,m-1} - 2u_{n,m}] + \omega_0^2 \sin u_{n,m} \quad (1.1)$$

J.M.Tamga *et al.* [39] studied how a weak initial uniform perturbation can evolve spontaneously into nonlinear localized modes with large amplitudes and investigated the solitary-wave and particle-like properties of these robust nonlinear entities.

The purpose of this paper is to show the existence of *uniform sliding states* [38] and *periodic travelling wave solutions* for 2d discrete models like (1.1) by using topological and variational methods as well. Variational methods are used in [2, 21, 29, 36] for 1d discrete FPU type lattice equations, while topological methods are applied in [26] to 1d damped discrete sine-Gordon lattice equations. Center manifold reduction methods are used in [15, 24, 25] to find travelling waves on nonlinear lattice equations. Dissipative lattice systems are investigated in [10]. The

existence and stability of solitary waves are studied also in [19, 20] for FPU lattices.

Our paper has the following structure: Section 2 discusses the mathematical formulation of the travelling wave solutions in two-dimensional lattices. The proof of uniform sliding states in equations like (1.1) is given in Section 3. There we also discuss their stability. In Section 4, using variational methods, we prove the existence of several types of periodic travelling wave solutions in equations like (1.1). Section 5 is devoted to multiplicity results of periodic travelling waves. Applications of our methods to more general nonlinear lattices are discussed in Section 6.

2 Formulation of the problem

In this paper, we consider the infinite system of ODEs

$$\ddot{u}_{n,m} = (\Delta u)_{n,m} - f(u_{n,m}), \quad (n, m) \in \mathbb{Z}^2 \quad (2.1)$$

on the two dimensional integer lattice $\mathbb{Z}^2$ under the following conditions for $f \in C^1(\mathbb{R}, \mathbb{R})$:

(H1) f is odd, i.e $f(-x) = -f(x) \forall x \in \mathbb{R}$.

(H2) f is 2π -periodic, i.e. $f(x + 2\pi) = f(x) \forall x \in \mathbb{R}$.

Δ denotes the discrete Laplacian defined as

$$(\Delta u)_{n,m} = u_{n+1,m} + u_{n-1,m} + u_{n,m+1} + u_{n,m-1} - 4u_{n,m} \quad (2.2)$$

We set

$$L := \max_{\mathbb{R}} |f'(x)|.$$

So f is globally Lipschitz continuous on $\mathbb{R}$, i.e. $|f(x) - f(y)| \leq L|x - y| \forall x, y \in \mathbb{R}$.

For $f(u) = \omega_0^2 \sin u$ and $G = 1$, we get the 2d discrete sine-Gordon lattice equation (1.1).

There is a wealth of structures that arise in problems on a two-dimensional lattice, stemming from the fact that one may consider different directions of motion of a travelling wave.

As a result of the symmetry imposed by the lattice $\mathbb{Z}^2$, the existence and speed of a wave generally will depend on the direction $e^{i\theta}$ of motion, with a special role of those directions for which the slope $\tan \theta$ is rational. Let $\theta \in \mathbb{R}$ be given, consider a solution of (2.1) of the form

$$u_{n,m}(t) = U(n \cos \theta + m \sin \theta - \nu t) \quad (2.3)$$

for some $\nu \in \mathbb{R}$ and $U : \mathbb{R} \rightarrow \mathbb{R}$. We may consider solutions of (2.3) to be travelling waves on the lattice $\mathbb{Z}^2$, in the direction $e^{i\theta}$. Substitution of (2.3) into (2.1) leads to the equation

$$\nu^2 U''(z) = U(z + \cos \theta) + U(z - \cos \theta) + U(z + \sin \theta) + U(z - \sin \theta) - 4U(z) - f(U(z)) \quad (2.4)$$

with $z = n \cos \theta + m \sin \theta - \nu t$.

If $\nu \neq 0$, equation (2.4) is a functional differential equation of a mixed type, where "mixed" refers to the fact that the equation involves both forward and backward translations of the argument of the solution U .

Note, that if θ is an integer multiple of $\pi/2$, then (2.4) reduces to the equation [15]

$$\nu^2 U''(z) = U(z+1) - 2U(z) + U(z-1) - f(U(z)) \quad (2.5)$$

On the other hand, if $\theta - \pi/4$ is an integer multiple of $\pi/2$, then under the change of variables $\psi(z) = U(2^{-1/2}z)$ equation (2.4) is equivalent to the equation

$$\nu^2 \psi''(z) = \psi(z+1) - 2\psi(z) + \psi(z-1) - 2^{-1}f(\psi(z)) \quad (2.6)$$

which is the same as (2.5), but with a rescaled function of $f(u)$.

For other values of θ , one does not expect any reduction of (2.4) to a simpler equation.

The limit $\nu \rightarrow 0$ in (2.4) results in a singular perturbation problem. Indeed, when $\nu = 0$ then (2.4) is in fact a functional-difference equation, not a functional-differential equation. In this case, the solution is defined on the set $\mathcal{Z} \subset \mathbb{R}$ given by

$$\mathcal{Z} = \{ z = n \cos \theta + m \sin \theta, (n, m) \in \mathbb{Z}^2 \}$$

The set $\mathcal{Z}$ is either a discrete subset or a countable dense subset of $\mathbb{R}$, depending on whether the quantity $\tan \theta$ is rational or irrational. Such functional-difference equations were studied in the series of papers [4, 5, 6]. We present a discussion in this direction in the point 4. of Section 6.

3 Uniform Sliding States

In this section, we prove the existence of *uniform sliding states* [38] in a two-dimensional lattice. In particular, we are looking for a solution $U(z)$ of (2.4) with the properties

$$U(z+T) = U(z) + 2\pi, \quad U(-z) = -U(z) \quad (3.1)$$

for some $T > 0$. $U(z)$ is called *the hull function* [38].

Theorem 1 *Let conditions (H1) and (H2) hold. Then equation (2.4) has a solution with (3.1) for any $\nu > 0$ and $T > 0$ satisfying the following condition:*

i) *either $\nu \geq 1$, or $0 < \nu < 1$ along with*

$$\frac{\pi}{r_{\nu,j}} k \neq T \quad \forall k \in \mathbb{N}, \quad j = 1, 2, \dots, n_\nu,$$

where $r_{\nu,j}$, $j = 1, 2, \dots, n_\nu$ are all positive solutions of the equation

$$\nu^2 = \frac{\sin^2(r_\nu \sin \theta) + \sin^2(r_\nu \cos \theta)}{r_\nu^2}.$$

Moreover, if $\nu > 1$ and $LT^2 < 4\pi^2(\nu^2 - 1)$ then this solution is unique.

Proof. We split $U(z)$ as follows

$$U(z) = \frac{2\pi}{T}z + x(z)$$

with $x(z+T) = x(z)$ and $x(-z) = -x(z)$. So we get

$$\begin{aligned} \nu^2 x''(z) &= x(z + \cos \theta) + x(z - \cos \theta) \\ &+ x(z + \sin \theta) + x(z - \sin \theta) - 4x(z) - f\left(\frac{2\pi}{T}z + x(z)\right). \end{aligned} \quad (3.2)$$

On a Hilbert space

$$X_1 := \left\{ x \in L_{loc}^2(\mathbb{R}, \mathbb{R}) \mid x(z+T) = x(z), \quad x(-z) = -x(z) \right\}$$

we define a linear mapping $L_1 : D(L_1) \subset X_1 \rightarrow X_1$ as follows

$$L_1(x)(z) := -\nu^2 x''(z) + x(z + \cos \theta) + x(z - \cos \theta) + x(z + \sin \theta) + x(z - \sin \theta) - 4x(z)$$

where $D(L_1) := X_1 \cap W_{loc}^{2,2}(\mathbb{R}, \mathbb{R})$. We also define the nonlinear operator $N_1 : X_1 \rightarrow X_1$ by

$$N_1(x)(z) := f\left(\frac{2\pi}{T}z + x(z)\right)$$

The Hilbert space X_1 is considered with the inner product

$$\langle x, y \rangle := \int_0^T x(z)y(z) dz$$

and the corresponding norm is denoted by $\|\cdot\|$. Then (3.2) has the form

$$L_1 x = N_1(x). \quad (3.3)$$

We want to show that the mapping L_1 is invertible and to derive an upper bound for the norm of its inverse.

According to Appendix, the spectrum $\sigma(L_1)$ is given by

$$\sigma(L_1) = \left\{ k^2 \nu^2 \frac{4\pi^2}{T^2} - 4 \sin^2\left(\frac{\pi}{T}k \sin \theta\right) - 4 \sin^2\left(\frac{\pi}{T}k \cos \theta\right) \mid k \in \mathbb{N} \right\}.$$

Let us put

$$g_\theta(r) = \frac{\sin^2(r \sin \theta) + \sin^2(r \cos \theta)}{r^2}.$$

Then

$$g_\theta(r) \leq \frac{\sin^2(r \sin \theta)}{r^2} + \frac{\sin^2(r \cos \theta)}{r^2} \leq 1.$$

So $g_\theta(r)$ attains its maximum 1 on $[0, \infty)$ at $r = 0$.

Hence if $\nu \geq 1$ then $0 \notin \sigma(L_1)$.

Since $g_\theta(r) \rightarrow 0$ as $r \rightarrow \infty$, for $0 < \nu < 1$, there is a finite number of positive $r_{\nu,j}$, $j = 1, 2, \dots, n_\nu$ solving

$$\nu^2 = g_\theta(r).$$

If assumption i) holds then again $0 \notin \sigma(L_1)$. Furthermore, there is an orthonormal basis $\{e_k\}_{k \in \mathbb{N}}$ of the corresponding eigenvectors

$$e_k := \left\{ \frac{1}{\sqrt{T}} \sin \frac{2\pi k}{T} z \right\}_{k \in \mathbb{N}}.$$

If $0 \notin \sigma(L_1)$ then for $x(z) = \sum_{k \in \mathbb{N}} c_k e_k$ we get

$$L_1^{-1}x = \sum_{k \in \mathbb{N}} \frac{c_k}{h_\theta(\pi k/T)} e_k$$

for

$$h_\theta(r) := 4 \left(\nu^2 r^2 - \sin^2(r \sin \theta) - \sin^2(r \cos \theta) \right).$$

This shows that the mapping $L_1^{-1} : X_1 \rightarrow X_1$ is compact [7, p. 13]. Next, the operator $N_1 : X_1 \rightarrow X_1$ is bounded and globally Lipschitz continuous with a constant L . Then (3.3) has the form

$$x = L_1^{-1} N_1(x). \quad (3.4)$$

Applying the Schauder fixed point theorem [7, p. 22] to (3.4) we obtain the existence part of Theorem 1.

Furthermore, for $\nu > 1$ we have

$$\|L_1^{-1}x\|^2 = \sum_{k \in \mathbb{N}} \frac{T^4}{16\pi^4 k^4} \frac{c_k^2}{(\nu^2 - g_\theta(\pi k/T))^2} \leq \sum_{k \in \mathbb{N}} \frac{T^4}{16\pi^4} \frac{c_k^2}{(\nu^2 - 1)^2} = \frac{T^4}{16\pi^4(\nu^2 - 1)^2} \|x\|^2.$$

Hence $\|L_1^{-1}\| \leq \frac{T^2}{4\pi^2(\nu^2 - 1)}$. If $\frac{T^2 L}{4\pi^2(\nu^2 - 1)} < 1$ then the Banach fixed point theorem [9, p. 24] gives the uniqueness result of Theorem 1. ■

Remark 1 The function $\tilde{U}(z) := U(-z)$ for $U(z)$ from Theorem 1 satisfies

$$\tilde{U}(z + T) = \tilde{U}(z) - 2\pi. \quad (3.5)$$

So we also get a solution of (2.4) satisfying (3.5).




Figure 1: The graphs of functions $\tilde{g}_{\pi/4}(\pi k/T)$ for $k = 1, 2, 3, 4, 5$ and $0 < T < 4$.

We note that condition i) is equivalent to $\nu \neq \tilde{g}_{\theta}(\pi k/T)$ for all $k \in \mathbb{N}$, and a given $T > 0$ with

$$\tilde{g}_{\theta}(r) := \frac{\sqrt{\sin^2(r \sin \theta) + \sin^2(r \cos \theta)}}{r}.$$

For a fixed $k \in \mathbb{N}$, the function $T \rightarrow \tilde{g}_{\theta}(\pi k/T)$ is oscillating near $T \sim 0$. So it is impossible to draw a complete diagram in the $\nu - T$ -plane where condition i) holds. We draw parts of graphs of these functions for $\theta = \pi/4$, $k = 1, 2, 3, 4, 5$ and $0 < T < 4$ in Figure 1. We can expect from Figure 1 the following result.

Corollary 1 *Let conditions (H1) and (H2) hold. If $\nu > T\sqrt{2}/\pi$ then equation (2.4) has a solution with (3.1). If $\nu > T\frac{\sqrt{L+8}}{2\pi}$ then this solution is unique.*

Proof. If $\nu > T\sqrt{2}/\pi$ then from $h_{\theta}(r) \geq 4\nu^2 r^2 - 8$, we get $\|L_1^{-1}\| \leq \frac{T^2}{4\nu^2 \pi^2 - 8T^2}$. This gives the existence part applying the Schauder fixed point theorem to (3.4). Next, $\nu > T\frac{\sqrt{L+8}}{2\pi}$ gives $\|L_1^{-1}\|L \leq \frac{LT^2}{4\nu^2 \pi^2 - 8T^2} < 1$. The use of the Banach fixed point theorem to (3.4) implies the uniqueness part. We note that the conditions of Corollary 1 implies condition i) of Theorem 1. ■

Now we improve Theorem 1 as follows:

Proposition 1 *Let conditions (H1), (H2) and i) hold. Then equation (2.4) has a solution $U(z)$ satisfying (3.1) which is accumulated by similar ones, i.e. there are sequences $\{U_n(z)\}_{n \in \mathbb{N}}$ and $\{T_n\}_{n \in \mathbb{N}}$ such that $U_n(z)$ is a solution of (2.4) satisfying (3.1) with T_n along with $T_n \rightarrow T$ and $U_n(z) \rightrightarrows U(z)$ uniformly on $[-2T, 2T]$ as $n \rightarrow \infty$.*

Proof. We know from the proof of Theorem 1 that all solutions of (3.3) belong into a ball $B_0 \subset X_1$. If there are infinitely many solutions of (3.3) then their accumulation point $x(z)$

gives $U(z) = \frac{2\pi}{T}z + x(z)$ which satisfies properties of Proposition 1. Here we use: $L_1^{-1} : X_1 \rightarrow D(L_1) \subset W_{loc}^{2,2}(\mathbb{R}, \mathbb{R}) \subset C(\mathbb{R}, \mathbb{R})$. If there are finitely many solutions $\{x_k\}_{k=1}^{M_0} \subset B_0$ of (3.3) then from the fact that the Leray-Schauder topological degree $\deg(I - L_1^{-1}N_1, B_0, 0) = 1$ (cf. [7, p. 57], [9, p. 65]), there is an x_{n_0} such that its local Leray-Schauder topological degree $\deg(I - L_1^{-1}N_1, x_{n_0}, 0) \neq 0$ (cf. [9, p. 69], [28, p. 192]). Next, slightly changing T , condition i) is still satisfied and $I - L_1^{-1}N_1$ depends continuously on T . This implies, that for any $\hat{T} \sim T$, there is a solution $\hat{U}(z) = \frac{2\pi}{\hat{T}}z + \hat{x}(z)$ of (2.4) satisfying (3.1) with $\hat{T}$ for $\hat{x}(z)$ near $x_{n_0}(z)$. Consequently, $U(z) = \frac{2\pi}{T}z + x_{n_0}(z)$ satisfies the properties of Proposition 1. The proof is finished. $\blacksquare$

Next we note that for $\theta = 0$ we get (2.5) and $g_0(r) = \frac{\sin^2 r}{r^2}$. Condition i) of Theorem 1 for $0 < v < 1$ and $T = 2\pi$ becomes

$$\nu \neq \frac{2|\sin k/2|}{k}, \quad \forall k \in \mathbb{N}.$$

In terms of [38, p. 1604], this means that condition i) of Theorem 1 is a *superharmonic nonresonance assumption*. In [38], hull functions were calculated numerically in the 1d FK model with a driving force and with damping. Theorem 1 predicts that these solutions persist as exact solutions in the absence of a driving force and damping and in 2d. Moreover, Proposition 1 asserts that under a superharmonic nonresonance there are continuum many uniform sliding states accumulated by sequences of uniform sliding states. Hence they are clearly neither asymptotically stable nor hyperbolic. To illustrate that how the stability of uniform sliding states is sophisticated, we consider

$$\nu^2 U''(z) = 2U\left(z + \frac{\sqrt{2}}{2}\right) + 2U\left(z - \frac{\sqrt{2}}{2}\right) - 4U(z) - \varepsilon \sin U(z) \quad (3.6)$$

with $\varepsilon \neq 0$ small. Taking $T = \sqrt{2}$, $1 \geq |\varepsilon| > 0$ and $\nu > \frac{3\sqrt{2}}{2\pi} \doteq 0.6752$, Corollary 1 ensures the existence of a unique odd $\sqrt{2}$ -periodic solution of

$$\nu^2 x''(z) = 2x\left(z + \frac{\sqrt{2}}{2}\right) + 2x\left(z - \frac{\sqrt{2}}{2}\right) - 4x(z) - \varepsilon \sin\left(\sqrt{2}\pi z + x(z)\right) \quad (3.7)$$

with $x(z) = O(\varepsilon)$. Then

$$u_{n,m}^0(t) := U\left(\frac{\sqrt{2}}{2}(n+m) - \nu t\right) = \pi(n+m) - \sqrt{2}\pi\nu t + x\left(\frac{\sqrt{2}}{2}(n+m) - \nu t\right)$$

satisfies $u_{n+2,m}^0(t) = u_{n,m+2}^0(t) = u_{n,m}^0(t) + 2\pi$. Hence we consider (2.1) under periodic conditions

$$u_{n+2,m}(t) = u_{n,m+2}(t) = u_{n,m}(t) + 2\pi. \quad (3.8)$$

Next we compute

$$\cos(u_{n,m}^0(t)) = \cos(\pi(n+m) - \sqrt{2}\pi\nu t) + O(\varepsilon) = \cos \pi(n+m) \cos \sqrt{2}\pi\nu t + O(\varepsilon).$$

Consequently, the linearization of (2.1) under conditions (3.8) at $u_{n,m}^0(t)$ up to $O(\varepsilon^2)$ -terms has the form

$$\begin{aligned}\ddot{v}_{0,0} &= 2v_{1,0} + 2v_{0,1} - 4v_{0,0} - \varepsilon \cos \sqrt{2}\pi \nu t v_{0,0} \\ \ddot{v}_{1,0} &= 2v_{0,0} + 2v_{1,1} - 4v_{1,0} + \varepsilon \cos \sqrt{2}\pi \nu t v_{1,0} \\ \ddot{v}_{0,1} &= 2v_{0,0} + 2v_{1,1} - 4v_{0,1} + \varepsilon \cos \sqrt{2}\pi \nu t v_{0,1} \\ \ddot{v}_{1,1} &= 2v_{0,1} + 2v_{1,0} - 4v_{1,1} - \varepsilon \cos \sqrt{2}\pi \nu t v_{1,1} .\end{aligned}\tag{3.9}$$

For $w := v_{0,0} - v_{1,1}$, (3.9) gives the Mathieu equation

$$\ddot{w} + \left(4 + \varepsilon \cos \sqrt{2}\pi \nu t\right) w = 0 .\tag{3.10}$$

For $\varepsilon > 0$ small, we get a resonance in (3.10) at $\sqrt{2}\pi \nu = 4/n$, $n \in \mathbb{N}$, [8]. This resonance is different from superharmonic one. Then from $\nu > \frac{3\sqrt{2}}{2\pi}$ we obtain $n = 1$, i.e. (3.10) is resonant only at $\nu = \nu_0 := 2\sqrt{2}/\pi \doteq 0.9003 > 0.6752$. Then according to [8], (3.10) is stable for $\nu \sim \nu_0$ in a narrow interval $\mathcal{I}$, and it is unstable for ν outside of $\mathcal{I}$. The smaller $\varepsilon > 0$ the narrow $\mathcal{I}$. So then (2.1) is also unstable at $u_{n,m}^0(t)$ for ν outside of $\mathcal{I}$. Of course, for a general θ , the analysis is much more difficult. So we do not carry out it in this paper.

Finally, sliding states of 1d FK models are investigated also in [17, 37].

4 Periodic Travelling Waves

First, in this Section, we are looking for a solution of (2.4) with the property

$$U(z + T) = -U(z) + 2\pi\tag{4.1}$$

for some $T > 0$. Of course then

$$U(z + 2T) = U(z) .\tag{4.2}$$

We note that conditions (H1) and (H2) imply

$$f(\pi - x) = -f(\pi + x) .\tag{4.3}$$

Hence $f(\pi) = 0$ and $x(z) = \pi$ is a trivial/constant solution of (2.4) satisfying (4.1).

We need the following definition and Theorem 2.8 from [14] (see also [28, p. 185, and p. 216, Exercise 3.]).

Definition 1 *Let H be a Hilbert space and $L : H \rightarrow H$ be a symmetric bounded linear operator. If the dimension of the subspace of all eigenvectors of L with negative eigenvalues is finite then it is called as the Morse index of L and it is denoted by $\text{index } L$.*

Theorem 2 *Let $\langle \cdot, \cdot \rangle$ be an inner product on a Hilbert space H with the corresponding norm $\|\cdot\|$. Let us consider a function $f \in C^2(H, \mathbb{R})$ of a form*

$$f(u) = \frac{1}{2} \langle Lu, u \rangle + g(u)$$

with the following properties:

- I) L is an isomorphism possessing the Morse index, i.e. the dimension of the subspace of all eigenvectors of L with negative eigenvalues is finite.
- II) $\nabla g : H \rightarrow H$ is a compact operator.
- III) There is a constant $M > 0$ such that $\|\nabla g(u)\| \leq M \forall u \in H$.
- IV) $\nabla f(0) = 0$ and the Morse index of $\text{Hess } f(0)$ exists, i.e. the dimension of the subspace of all eigenvectors of $\text{Hess } f(0)$ with negative eigenvalues is finite.

If the following condition is satisfied:

$$\text{index } L \notin [\text{index } \text{Hess } f(0), \text{index } \text{Hess } f(0) + \dim \ker \text{Hess } f(0)]$$

then f possesses a nonzero critical point on H .

Remark 2 When $L = \alpha I + B$ for a constant $\alpha > 0$ and a compact linear symmetric operator $B : H \rightarrow H$, then the both linear operators L and $\text{Hess } f(0)$ have Morse indices, i.e. the dimensions of the subspaces of all eigenvectors of L and $\text{Hess } f(0)$ with negative eigenvalues are finite, respectively (cf. [7, p. 75]).

We denote by $(a, b) \subset \mathbb{R}$ the interval with end points a, b for either $a < b$ or $b < a$. Now we can prove the following result.

Theorem 3 Let conditions (H1), (H2) hold and $f'(\pi) \neq 0$. Moreover, suppose

- a) either $\nu \geq 1$, or $0 < \nu < 1$ and

$$\frac{\pi}{2r_{\nu,j}}(2k-1) \neq T \quad \forall k \in \mathbb{N}, \quad j = 1, 2, \dots, n_\nu,$$

where $r_{\nu,j}$, $j = 1, 2, \dots, n_\nu$ are defined in Theorem 1.

Furthermore, we assume

- b) There exists a $k_0 \in \mathbb{N}$ such that

$$\nu^2 \in \left(g_\theta \left(\frac{\pi}{2T} (2k_0 - 1) \right), g_\theta \left(\frac{\pi}{2T} (2k_0 - 1) \right) + \frac{T^2}{\pi^2 (2k_0 - 1)^2} f'(\pi) \right).$$

Then (2.4) has a nonconstant solution satisfying (4.1).

Proof. We split $U(z)$ as follows

$$U(z) = \pi + x(z)$$

with $x(z + T) = -x(z)$. So we get

$$\begin{aligned} \nu^2 x''(z) &= x(z + \cos \theta) + x(z - \cos \theta) \\ &+ x(z + \sin \theta) + x(z - \sin \theta) - 4x(z) - f(\pi + x(z)). \end{aligned} \quad (4.4)$$

We take Hilbert spaces

$$\begin{aligned} X_2 &:= \left\{ x \in L_{loc}^2(\mathbb{R}, \mathbb{R}) \mid x(z + T) = -x(z) \right\} \\ Y_2 &:= \left\{ x \in W_{loc}^{1,2}(\mathbb{R}, \mathbb{R}) \mid x(z + T) = -x(z) \right\} \end{aligned}$$

with the above inner product $\langle x, y \rangle$ and

$$(x, y) := \int_0^T x'(z) y'(z) dz,$$

respectively. Moreover, we consider operators

$$\begin{aligned} L_2 x &:= x(z + \cos \theta) + x(z - \cos \theta) \\ &+ x(z + \sin \theta) + x(z - \sin \theta) - 4x(z) \\ N_2(x) &:= -f(\pi + x(z)). \end{aligned}$$

On Y_2 we set the function

$$H(x) := \int_0^T \left\{ \nu^2 \frac{x'(z)^2}{2} + (x(z + \cos \theta) + x(z + \sin \theta))x(z) - 2x(z)^2 - F(\pi + x(z)) \right\} dz$$

for $F(x) = \int_0^x f(s) ds$. Note $H \in C^2(Y_2, \mathbb{R})$ and

$$\begin{aligned} (DH(x), v) &= \int_0^T \left\{ \nu^2 x'(z) v'(z) + (x(z + \cos \theta) + x(z + \sin \theta))v(z) \right. \\ &\quad \left. + (v(z + \cos \theta) + v(z + \sin \theta))x(z) - 4x(z)v(z) - f(\pi + x(z))v(z) \right\} dz \end{aligned}$$

If x is a critical point of H , then, for every $v \in Y_2$,

$$\begin{aligned} 0 &= (DH(x), v) = \int_0^T \left\{ \nu^2 x'(z) v'(z) + (x(z + \cos \theta) + x(z + \sin \theta))v(z) \right. \\ &\quad \left. + (v(z + \cos \theta) + v(z + \sin \theta))x(z) - 4x(z)v(z) - f(\pi + x(z))v(z) \right\} dz \end{aligned} \quad (4.5)$$

with $x \in Y_2$. But for any c , we have

$$\int_0^T x(z + c)v(z) dz = \int_c^{T+c} x(s)v(s - c) ds = \int_0^T x(s)v(s - c) ds$$

since $x, v \in Y_2$, gives the T -periodicity of $s \rightarrow x(s)v(s - c)$. Consequently, (4.5) gives us $x \in Y_2 \cap W_{loc}^{2,2}(\mathbb{R}, \mathbb{R})$ along with

$$0 = \int_0^T \left\{ -\nu^2 x''(z) + x(z + \cos \theta) + x(z + \sin \theta) + x(z - \cos \theta) + x(z - \sin \theta) - 4x(z) - f(\pi + x(z)) \right\} v(z) dz \quad (4.6)$$

Since, (4.6) holds for any $v \in Y_2$, we see that the critical points of H are solutions of (4.4).

Next, the Riesz representation theorem of bounded linear functionals on Hilbert spaces [33] gives a linear mapping $K : X_2 \rightarrow Y_2$ defined by the equality

$$(Kh, y) = \langle h, y \rangle \quad \forall y \in Y_2.$$

Then

$$\begin{aligned} \nabla H(x) &= \nu^2 x + KL_2 x + KN_2(x), \\ \text{Hess } H(x) &= \nu^2 I + KL_2 + KDN_2(x), \end{aligned}$$

where we consider

$$L_2, N_2 : Y_2 \hookrightarrow X_2 \rightarrow X_2.$$

Hence

$$\tilde{L}_2 := KL_2, \quad \tilde{N}_2(x) := KN_2(x)$$

are compact operators. Moreover, $\tilde{N}_2 : Y_2 \rightarrow Y_2$ is bounded.

Since by (4.3), $f(\pi) = 0$, equation (4.4) has a trivial solution $x(z) = 0$. We show the existence of a nontrivial/nonzero one by using Theorem 2. The boundedness of $\tilde{N}_2$ gives that $\nabla H(x)$ is asymptotically linear at infinity with an asymptote $\bar{L}_2 = \nu^2 I + \tilde{L}_2$. Next using an orthogonal basis

$$\left\{ \cos \frac{\pi}{T}(2k-1)z, \sin \frac{\pi}{T}(2k-1)z \right\}_{k \in \mathbb{N}}$$

we see that

$$\begin{aligned} \sigma(\bar{L}_2) &= \left\{ \nu^2 - g_\theta \left(\frac{\pi}{2T}(2k-1) \right) \right\}_{k \in \mathbb{N}}, \\ \sigma(\text{Hess } H(0)) &= \left\{ \nu^2 - g_\theta \left(\frac{\pi}{2T}(2k-1) \right) - \frac{T^2}{\pi^2(2k-1)^2} f'(\pi) \right\}_{k \in \mathbb{N}}. \end{aligned}$$

Assumption a) gives $\ker \bar{L}_2 = \{0\}$. According to Definition 1, the Morse indices of $\bar{L}_2$ and $\text{Hess } H(0)$ are given by

$$\begin{aligned} \text{index } \bar{L}_2 &= \# \left\{ k \in \mathbb{N} \mid \nu^2 < g_\theta \left(\frac{\pi}{2T}(2k-1) \right) \right\} \times 2, \\ \text{index Hess } H(0) &= \\ &= \# \left\{ k \in \mathbb{N} \mid \nu^2 < g_\theta \left(\frac{\pi}{2T}(2k-1) \right) + \frac{T^2}{\pi^2(2k-1)^2} f'(\pi) \right\} \times 2. \\ \dim \ker \text{Hess } H(0) &= \\ &= \# \left\{ k \in \mathbb{N} \mid \nu^2 = g_\theta \left(\frac{\pi}{2T}(2k-1) \right) + \frac{T^2}{\pi^2(2k-1)^2} f'(\pi) \right\} \times 2. \end{aligned}$$

Hence condition b) implies

$$\text{index } \bar{L}_2 \notin \left[\text{index Hess } H(0), \text{index Hess } H(0) + \dim \ker \text{Hess } H(0) \right].$$

Then Theorem 2 ensures the existence of a nonzero critical point of H . ■

Similarly we have the next results.

Theorem 4 *Let conditions (H1), (H2) hold and $f'(0) \neq 0$. Moreover, suppose condition a) of Theorem 3 along with*

c) *There exists a $k_0 \in \mathbb{N}$ such that*

$$\nu^2 \in \left(g_\theta \left(\frac{\pi}{2T} (2k_0 - 1) \right), g_\theta \left(\frac{\pi}{2T} (2k_0 - 1) \right) + \frac{T^2}{\pi^2 (2k_0 - 1)^2} f'(0) \right).$$

Then (2.4) has a nonzero solution satisfying

$$U(z + T) = -U(z). \quad (4.7)$$

Proof. We follow the proof of Theorem 3 for (2.4) in place of (4.4). So on Y_2 we set the function

$$\bar{H}(x) := \int_0^T \left\{ \nu^2 \frac{x'(z)^2}{2} + (x(z + \cos \theta) + x(z + \sin \theta))x(z) - 2x(z)^2 - F(x(z)) \right\} dz$$

Then,

$$\begin{aligned} \nabla \bar{H}(x) &= \nu^2 x + K L_2 x + K N_3(x), \\ \text{Hess } \bar{H}(x) &= \nu^2 I + K_2 L_2 + K D N_3(x), \end{aligned}$$

for

$$N_3(x) := -f(x(z)).$$

Consequently, we get

$$\begin{aligned} \text{index Hess } \bar{H}(0) &= \# \left\{ k \in \mathbb{N} \mid \nu^2 < g_\theta \left(\frac{\pi}{2T} (2k - 1) \right) + \frac{T^2}{\pi^2 (2k - 1)^2} f'(0) \right\} \times 2, \\ \dim \ker \text{Hess } \bar{H}(0) &= \# \left\{ k \in \mathbb{N} \mid \nu^2 = g_\theta \left(\frac{\pi}{2T} (2k - 1) \right) + \frac{T^2}{\pi^2 (2k - 1)^2} f'(0) \right\} \times 2. \end{aligned}$$

Hence condition c) implies

$$\text{index } \bar{L}_2 \notin \left[\text{index Hess } \bar{H}(0), \text{index Hess } \bar{H}(0) + \dim \ker \text{Hess } \bar{H}(0) \right].$$

Assumption a) of Theorem 3 gives $\ker \bar{L}_2 = \{0\}$. Then, Theorem 2 ensures the existence of a nonzero critical point of $\bar{H}$. ■

Remark 3 *Conditions (4.1) and (4.7) imply condition (4.2). But they give two different solutions of (2.4) satisfying (4.2).*

Theorem 5 *Let conditions (H1), (H2) hold and $f'(0) \neq 0$. Moreover, suppose condition i) of Theorem 1 along with*

d) There exists a $k_0 \in \mathbb{N}$ such that

$$\nu^2 \in \left(g_\theta \left(\frac{\pi}{T} k_0 \right), g_\theta \left(\frac{\pi}{T} k_0 \right) + \frac{T^2}{4\pi^2 k_0^2} f'(0) \right).$$

Then (2.4) has a nonzero solution satisfying

$$U(z+T) = U(z), \quad U(-z) = -U(z). \quad (4.8)$$

Proof. We consider (2.4) on the Hilbert spaces X_1 and

$$Y_1 := \left\{ x \in W_{loc}^{1,2}(\mathbb{R}, \mathbb{R}) \mid x(z+T) = x(z), \quad x(-z) = -x(z) \right\},$$

where Y_1 is endowed with the above inner product $(\cdot, \cdot)$.

Next, on Y_1 we take the above function $\bar{H}$. The rest of the proof is a modification of Theorem 3, but we present it here for completeness. Again, the Riesz representation theorem gives a linear mapping $\bar{K} : X_1 \rightarrow Y_1$ defined by the equality

$$(\bar{K}h, y) = \langle h, y \rangle \quad \forall y \in Y_1.$$

So

$$\begin{aligned} \nabla \bar{H}(x) &= \nu^2 x + \bar{K}L_2 x + \bar{K}N_3(x), \\ \text{Hess } \bar{H}(x) &= \nu^2 I + \bar{K}L_2 + \bar{K}DN_3(x), \end{aligned}$$

where we consider

$$L_2, N_3 : Y_1 \hookrightarrow X_1 \rightarrow X_1.$$

Clearly

$$\hat{L}_2 := \bar{K}L_2, \quad \hat{N}_3(x) := \bar{K}N_3(x)$$

are compact operators. Moreover, $\nabla \bar{H}(x)$ is asymptotically linear at infinity with an asymptote $\check{L}_2 = \nu^2 I + \hat{L}_2$. Using the orthogonal basis $\{e_k\}_{k \in \mathbb{N}}$, we get

$$\begin{aligned} \text{index } \check{L}_2 &= \# \left\{ k \in \mathbb{N} \mid \nu^2 < g_\theta \left(\frac{\pi}{T} k \right) \right\}, \\ \text{index Hess } \bar{H}(0) &= \# \left\{ k \in \mathbb{N} \mid \nu^2 < g_\theta \left(\frac{\pi}{T} k \right) + \frac{T^2}{4\pi^2 k^2} f'(0) \right\}, \\ \dim \ker \text{Hess } \bar{H}(0) &= \# \left\{ k \in \mathbb{N} \mid \nu^2 = g_\theta \left(\frac{\pi}{T} k \right) + \frac{T^2}{4\pi^2 k^2} f'(0) \right\}. \end{aligned}$$

Hence condition d) implies

$$\text{index } \check{L}_2 \notin \left[\text{index Hess } \bar{H}(0), \text{index Hess } \bar{H}(0) + \dim \ker \text{Hess } \bar{H}(0) \right].$$

Assumption i) of Theorem 1 gives $\ker \check{L}_2 = \{0\}$. Then Theorem 2 ensures the existence of a nonzero critical point of $\bar{H}$. ■

Applying the above results, for instance, we have the following

Corollary 2 *Let conditions (H1) and (H2) hold. If $\nu \geq 1$ and*

$$f'(0) > \frac{\pi^2}{T^2} \left(\nu^2 - g_\theta \left(\frac{\pi}{2T} \right) \right)$$

then (2.4) has a nonzero solution satisfying (4.7).

Corollary 3 *Let conditions (H1) and (H2) hold. If $\nu \geq 1$ and*

$$f'(0) > \frac{4\pi^2}{T^2} \left(\nu^2 - g_\theta \left(\frac{\pi}{T} \right) \right)$$

then (2.4) has a nonzero odd T -periodic solution.

Now let $\theta = \pi/4$. Then there is an $\nu_1 \in (0, 1)$ such that for any $\nu_1 < \nu < 1$, the equation

$$\nu^2 = g_{\pi/4}(r) = 2 \frac{\sin^2 \left(\frac{\sqrt{2}}{2} r \right)}{r^2}$$

has a unique positive solution r_ν . An approximate value of ν_1 is 0.2172, and $r_\nu \in (0, 3.6112)$. Taking step by step $k_0 = 2, 1, 1$ in Theorems 3, 4 and 5, we can derive from these theorems the following result.

Corollary 4 *Let $\theta = \pi/4$ and $\nu \in (\nu_1, 1)$. If*

$$\begin{aligned} \frac{\pi}{r_\nu} &< T < \frac{3\pi}{2r_\nu}, \\ f'(\pi) &> \left(\nu^2 - g_{\pi/4} \left(\frac{3\pi}{2T} \right) \right) \frac{9\pi^2}{T^2}, \\ f'(0) &< \frac{\pi^2}{T^2} \min \left\{ \nu^2 - g_{\pi/4} \left(\frac{\pi}{2T} \right), 4 \left(\nu^2 - g_{\pi/4} \left(\frac{\pi}{T} \right) \right) \right\}. \end{aligned} \tag{4.9}$$

Then (2.4) has 4 nontrivial/nonconstant solutions satisfying either (3.1), or (4.1), or (4.7), or (4.8), respectively.

On the other hand, taking $k_0 = 2$ in Theorem 4 we get

Corollary 5 *Let $\theta = \pi/4$ and $\nu \in (\nu_1, 1)$. If*

$$\begin{aligned} \frac{\pi}{r_\nu} &< T < \frac{3\pi}{2r_\nu}, \\ f'(0) &> \left(\nu^2 - g_{\pi/4} \left(\frac{3\pi}{2T} \right) \right) \frac{9\pi^2}{T^2}. \end{aligned} \tag{4.10}$$

Then (2.4) has a nonzero solution satisfying (4.7).

To be more concrete, we take $\theta = \pi/4$ and $\nu = 1/2$. Then we can numerically verify that approximately $r_{1/2} \doteq 2.6806$ and then the conditions of (4.9) hold for

$$1.17196 < T < 1.7579, \quad f'(0) < -5, \quad f'(\pi) > 16,$$

while the conditions of (4.10) hold for

$$1.17196 < T < 1.7579, \quad f'(0) > 16.$$

So we obtain the following result.

Corollary 6 *For any $\omega > 16$ and $1.17196 < T < 1.7579$, the 2d discrete sine-Gordon equation*

$$\ddot{u}_{n,m} - u_{n+1,m} - u_{n-1,m} - u_{n,m+1} - u_{n,m-1} + 4u_{n,m} + \omega \sin u_{n,m} = 0 \quad (4.11)$$

possesses 4 nontrivial/nonconstant travelling wave solutions of the form

$$u_{n,m}(t) = \pi + U\left(\frac{1}{\sqrt{2}}\left(n + m\right) - \frac{1}{2}t\right)$$

for $U(z)$ satisfying either (3.1), or (4.1), or (4.7), or (4.8), respectively.

Proof. First we change $u = w + \pi$. Then (4.11) becomes

$$\ddot{w}_{n,m} - w_{n+1,m} - w_{n-1,m} - w_{n,m+1} - w_{n,m-1} + 4w_{n,m} - \omega \sin w_{n,m} = 0. \quad (4.12)$$

Now we take $f(w) = -\omega \sin w$. Then $f'(0) = -\omega < 0$ and $f'(\pi) = \omega > 0$. The results follow from the above observation. The proof is finished. $\blacksquare$

We note that the solution $\bar{U}(z) := \pi + U(z)$ of (4.11) corresponding to condition (4.1) in Corollary 6 satisfies $\bar{U}(z + T) = -\bar{U}(z) + 4\pi$. So $\bar{U}(z + T) - 2\pi = -(\bar{U}(z) - 2\pi)$, and $\tilde{V}(z) = \bar{U}(z) - 2\pi$ determines a $2T$ -periodic travelling wave for (4.11) satisfying (4.7).

Of course, more layers of continuum many uniform sliding states and periodic travelling waves can be shown than in Corollaries 4, 5 and 6 by choosing different k_0 and θ in Theorems 3, 4 and 5.

5 Multiplicity Results

It is not difficult to see from [27], that $U(z) = 0$ is the only odd T -periodic solution of (2.4) when $(-L, L) \cap \sigma(L_1) = \emptyset$. On the other, we do not know in general about uniqueness or multiplicity of nonconstant/nonzero solutions in Theorems 3, 4 and 5. But it seems that enforcing assumptions, we could get rather multiplicity results of nonconstant/nonzero solutions than uniqueness. As an illustration, now we extend Theorem 5 as follows.

Theorem 6 *Let assumptions (H1), (H2) hold, $\nu > 1$ and $f'(0) > 0$. Moreover, suppose*

$$\mathbf{e1)} \quad \nu^2 \neq g_\theta\left(\frac{\pi k}{T}\right) + \frac{T^2}{4\pi^2 k^2} f'(0) \quad \forall k \in \mathbb{N},$$

e2) $\#\left\{k \in \mathbb{N} \mid \nu^2 < g_\theta\left(\frac{\pi}{T}k\right) + \frac{T^2}{4\pi^2 k^2}f'(0)\right\} \geq \left\lfloor \frac{T\sqrt{L}}{2\pi\sqrt{\nu^2-1}} \right\rfloor \geq 2$, where $[\cdot]$ is the integer part function.

Then (2.4) has at least 2 nonzero odd T -periodic solutions.

Proof. We work in the framework of Theorem 5. First, clearly Theorem 5 can be applied. So under the conditions of Theorem 6, we have an odd nonzero T -periodic solution $x_0(z)$ of (2.4). In order to show another one, we follow the proof of [28, Theorem 9.2], i.e. we must prove

$$|\text{index } \check{L}_2 - \text{index Hess } \bar{H}(0)| \geq \dim \ker \text{Hess } \bar{H}(x_0) \geq 2. \quad (5.1)$$

Since $\nu > 1$, $\text{index } \check{L}_2 = 0$. Next, e1) implies $\dim \ker \text{Hess } \bar{H}(0) = 0$, and we know

$$\text{index Hess } \bar{H}(0) = \#\left\{k \in \mathbb{N} \mid \nu^2 < g_\theta\left(\frac{\pi}{T}k\right) + \frac{T^2}{4\pi^2 k^2}f'(0)\right\}.$$

Next, we note that $\ker \text{Hess } \bar{H}(x_0)$ is determined by the equation

$$\nu^2 x''(z) = x(z + \cos \theta) + x(z - \cos \theta) + x(z + \sin \theta) + x(z - \sin \theta) - 4x(z) - f'(x_0(z))x(z). \quad (5.2)$$

We have $\dim \ker \text{Hess } \bar{H}(x_0) \leq \left\lfloor \frac{T\sqrt{L}}{2\pi\sqrt{\nu^2-1}} \right\rfloor$ from Lemma 1 below. Hence (5.1) follows from Lemma 1 and assumption e2). The proof is finished. $\blacksquare$

Lemma 1 *The dimension of all odd T -periodic solutions of (5.2) is less or equals to $\left\lfloor \frac{T\sqrt{L}}{2\pi\sqrt{\nu^2-1}} \right\rfloor$.*

Proof. Let $k_0 = \left\lfloor \frac{T\sqrt{L}}{2\pi\sqrt{\nu^2-1}} \right\rfloor + 1$. Split $X_1 = X_{k_0} \oplus X_{k_0}^\perp$ with $X_{k_0} := \text{span}\{e_k\}_{k=1}^{k_0-1}$. Let $P_{k_0} : X_1 \rightarrow X_1$ be the orthogonal projection onto X_{k_0} . Next, for $x = x_1 + x_2$, $x_1 \in X_{k_0}$, $x_2 \in X_{k_0}^\perp$ from (5.2) we obtain

$$\begin{aligned} \nu^2 x_1''(z) = & x_1(z + \cos \theta) + x_1(z - \cos \theta) + x_1(z + \sin \theta) + x_1(z - \sin \theta) - 4x_1(z) \\ & - P_{k_0} f'(x_0(z))(x_1(z) + x_2(z)), \end{aligned} \quad (5.3)$$

$$\begin{aligned} \nu^2 x_2''(z) = & x_2(z + \cos \theta) + x_2(z - \cos \theta) + x_2(z + \sin \theta) + x_2(z - \sin \theta) - 4x_2(z) \\ & - (I - P_{k_0}) f'(x_0(z))(x_1(z) + x_2(z)). \end{aligned} \quad (5.4)$$

Let

$$L_{k_0} := -\nu^2 x_2''(z) + x_2(z + \cos \theta) + x_2(z - \cos \theta) + x_2(z + \sin \theta) + x_2(z - \sin \theta) - 4x_2(z)$$

with $D(L_{k_0}) = X_{k_0}^\perp \cap W_{loc}^{2,2}(\mathbb{R}, \mathbb{R})$. Now, like in the proof of Theorem 1 we derive

$$\|L_{k_0}^{-1}\| \leq \frac{T^2}{4\pi^2 k_0^2 (\nu^2 - 1)}.$$

Consequently, (5.4) has the form

$$x_2 - L_{k_0}^{-1}(I - P_{k_0})f'(x_0)x_2 = L_{k_0}^{-1}(I - P_{k_0})f'(x_0)x_1. \quad (5.5)$$




Figure 2: The graph of (5.6).

Since

$$\|L_{k_0}^{-1}(I - P_{k_0})f'(x_0)x_2\| \leq \|L_{k_0}^{-1}\| \|I - P_{k_0}\| \|f'(x_0)x_2\| \leq \frac{T^2 L}{4\pi^2 k_0^2 (\nu^2 - 1)} \|x_2\|,$$

and $\frac{T\sqrt{L}}{2\pi k_0 \sqrt{\nu^2 - 1}} < 1$, the Banach fixed point theorem gives a unique solution $x_2 = x_2(x_1)$ of (5.5). Summarizing, the solvability of (5.2) is reduced to (5.3) with $x_2 = x_2(x_1)$. Since $\dim X_{k_0} = \left\lceil \frac{T\sqrt{L}}{2\pi\sqrt{\nu^2 - 1}} \right\rceil$, the proof is finished. $\blacksquare$

To show that assumptions e1) and e2) are consistent, we take $f(x) = \sin x$, so $f'(0) = L = 1$. Next, we consider $\theta = \pi/4$, $\nu = 2$ and solve the equation $g(r) = 0$ for a function

$$g(r) := \frac{2 \sin^2 \left(r \frac{\sqrt{2}}{2} \right)}{r^2} + \frac{1}{4r^2} - 4, \quad (5.6)$$

which has a unique solution r_0 (see Figure 2) with an approximative value 0.2880.

Hence e1) is equivalent to

$$\frac{\pi}{T}k \neq r_0, \quad \forall k \in \mathbb{N}, \quad (5.7)$$

while e2), when e1) holds, is equivalent to

$$\left\lceil \frac{T}{\pi} r_0 \right\rceil \geq \left\lceil \frac{T}{\pi} \frac{1}{2\sqrt{3}} \right\rceil \geq 2. \quad (5.8)$$

We note $\frac{1}{2\sqrt{3}} \doteq 0.2887 > r_0$. So the condition (5.8) holds only for bounded T/π , i.e. T can not be arbitrarily large, and $\left\lceil \frac{T}{\pi} r_0 \right\rceil \leq \left\lceil \frac{T}{\pi} \frac{1}{2\sqrt{3}} \right\rceil$. So it must be $\left\lceil \frac{T}{\pi} r_0 \right\rceil = \left\lceil \frac{T}{\pi} \frac{1}{2\sqrt{3}} \right\rceil$, which gives, according to e1), $k < \frac{T}{\pi} r_0 < \frac{T}{2\sqrt{3}\pi} < k+1$ for a $k \in N$, $k \geq 2$. That is: $\frac{\pi}{r_0} k < T < 2\sqrt{3}\pi(k+1)$. This implies $\frac{\pi}{r_0} k < 2\sqrt{3}\pi(k+1)$, i.e. $k < \frac{2\sqrt{3}r_0}{1-2\sqrt{3}r_0} \doteq 436.8855$. Hence the condition (5.8) is satisfied on the set $\bigcup_{k=2}^{436} \left(\frac{\pi}{r_0} k, 2\sqrt{3}\pi(k+1) \right)$ of disjunct intervals. We note that the first interval with $k = 2$ is approximately $(21.8154, 32.6484)$ and the last one with $k = 436$ is approximately $(4755.7599, 4755.7819)$. Summarizing we arrive at the following

Corollary 7 *For $f(x) = \sin x$, $\nu = 2$, $\theta = \pi/4$ and $T \in \bigcup_{k=2}^{436} \left(\frac{\pi}{r_0} k, 2\sqrt{3}\pi(k+1) \right)$, equation (2.4) has at least 2 nonzero odd T -periodic solutions.*

Other multiplicity result can be derived also as follows. We assume in the above results of Section 4 *nonresonance at infinity*, i.e. conditions i), a) of Theorems 1 and 3, respectively. Now we study an opposite case. So let us again consider (3.6) with $\varepsilon \neq 0$ small. We suppose $\nu_1 < \nu < 1$ and take

$$T = \pi/r_\nu.$$

Then (3.6) has the form

$$\nu^2 x + \hat{L}_2 x + \varepsilon \hat{N}_3(x) = 0, \quad x \in Y_1, \quad (5.9)$$

in the framework of Theorem 5 with $f(x) = \sin x$ and

$$\ker(\nu^2 I + \hat{L}_2) = \text{im}(\nu^2 I + \hat{L}_2)^\perp = \text{span} \{ \sin 2r_\nu z \}.$$

Hence (3.6) is *resonant at infinity*, i.e. it is *superharmonic resonant*. Next we split $x(z) = c \sin 2r_\nu z + x_1(z)$, $x_1 \in \text{im}(\nu^2 I + \hat{L}_2)$ to get from (5.9)

$$\nu^2 x_1 + \hat{L}_2 x_1 + \varepsilon P_1 \hat{N}_3(c \sin 2r_\nu z + x_1(z)) = 0, \quad (5.10)$$

$$\int_0^{\pi/r_\nu} \sin(c \sin 2r_\nu z + x_1(z)) \sin 2r_\nu z \, dz = 0, \quad (5.11)$$

where $P_1 : Y_1 \rightarrow Y_1$ is the orthogonal projection onto $\text{im}(\nu^2 I + \hat{L}_2)$. Using the implicit function theorem [9, p. 26], we can solve (5.10) to get $x_1(\varepsilon, z)$ for ε small with $x_1(0, z) = 0$. Inserting this solution into (5.11), we get the bifurcation equation

$$Q(\varepsilon, c) := \int_0^{\pi/r_\nu} \sin(c \sin 2r_\nu z + x_1(\varepsilon, z)) \sin 2r_\nu z \, dz = 0. \quad (5.12)$$

Clearly $Q(0, c)$ is an entire function on the complex plane $\mathbb{C}$ and

$$Q(0, c) := \frac{\partial}{\partial c} \tilde{Q}(c)$$

for

$$\begin{aligned}\tilde{Q}(c) &:= - \int_0^{\pi/r_\nu} \cos(c \sin 2r_\nu z) dz = \sum_{n=0}^{\infty} \int_0^{\pi/r_\nu} \frac{\sin^{2n} 2r_\nu z}{2n!} (-1)^{n+1} dz \cdot c^{2n} \\ &= \frac{\pi}{r_\nu} \sum_{n=0}^{\infty} \frac{1.3.5 \dots (2n-1)}{2.4.6 \dots 2n} \frac{1}{2n!} (-1)^{n+1} c^{2n} = \frac{\pi}{r_\nu} \sum_{n=0}^{\infty} \frac{1}{4^n n!^2} (-1)^{n+1} c^{2n}.\end{aligned}$$

Hence

$$\tilde{Q}(\iota\sqrt{c}) = -\frac{\pi}{r_\nu} \sum_{n=0}^{\infty} \frac{\phi(n)}{n!} c^n$$

for $c \in \mathbb{C}$ and the entire function

$$\phi(c) = \frac{1}{4^c \Gamma(c+1)}$$

which has only negative roots $-1, -2, \dots$. Using arguments of [40, Section 8], we see that the entire function $\sum_{n=0}^{\infty} \frac{\phi(n)}{n!} c^n$ has only real and negative roots on $\mathbb{C}$. So $\tilde{Q}(c)$ has only real roots on $\mathbb{C}$ and they are infinitely many. Then again by [40], $Q(0, c)$ has also only real roots on $\mathbb{C}$ separated by roots of $\tilde{Q}(c)$. This implies that there are infinitely many real roots of $Q(0, c)$ with odd multiplicity. Indeed, if it could exist c_0 such that $Q(0, c)$ would have on (c_0, ∞) only roots with even multiplicity then $Q(0, c)$ would not change the sign on (c_0, ∞) . So $\tilde{Q}(c)$ would be monotonic on this interval. This would contradict to that $\tilde{Q}(c)$ has infinitely many roots on this interval. Here we use the evenness of this function. Next, each real root of $Q(0, c)$ with odd multiplicity has a nonzero local Brouwer topological degree [9, p. 69], which implies that $Q(\varepsilon, c)$ has infinitely many real roots, so it is oscillatory on $\mathbb{R}$. Summarizing we arrive at the following

Theorem 7 *Let $\nu_1 < \nu < 1$. Then (3.6) has infinitely many odd π/r_ν -periodic solutions $\{U_n(z)\}_{n \in \mathbb{N}}$ with*

$$|U_n(z) - c_n \sin 2r_\nu z| \leq \tilde{K}|\varepsilon|$$

for $c_n \rightarrow \infty$ as $n \rightarrow \infty$ and a constant $\tilde{K} > 0$.

Theorem 7 states that under superharmonic resonance $T = \pi/r_\nu$, (3.6) possesses infinitely many T -periodic travelling waves. Note $r_\nu \in (0, 3.6112)$.

6 Concluding Remarks

1.

The same approach as in Section 4, we can apply to

$$\begin{aligned}\ddot{u}_{n,m} &= g(u_{n+1,m} - u_{n,m}) - g(u_{n,m} - u_{n-1,m}) \\ &\quad + g(u_{n,m+1} - u_{n,m}) - g(u_{n,m} - u_{n,m-1}) - f(u_{n,m}),\end{aligned}\tag{6.1}$$

where $f, g \in C^1(\mathbb{R}, \mathbb{R})$ are bounded odd. Then we study

$$\begin{aligned} \nu^2 U''(z) = & g(U(z + \cos \theta) - U(z)) - g(U(z) - U(z - \cos \theta)) \\ & + g(U(z + \sin \theta) - U(z)) - g(U(z) - U(z - \sin \theta)) - f(U(z)). \end{aligned} \quad (6.2)$$

Following the proof of Theorem 5 we have

Theorem 8 *Assuming $f, g \in C^1(\mathbb{R}, \mathbb{R})$ are bounded odd along with*

$$4\pi^2 \nu^2 - 4\pi^2 g'(0) g_\theta \left(\frac{\pi}{T} \right) < T^2 f'(0), \quad (6.3)$$

then (6.2) possesses a nonzero odd T -periodic solution.

Proof. This result follows like in Theorem 5 for the function

$$\tilde{H}(x) := \int_0^T \left\{ \nu^2 \frac{x'(z)^2}{2} - G(x(z + \cos \theta) - x(z)) - G(x(z + \sin \theta) - x(z)) - F(x(z)) \right\} dz$$

with $G(x) := \int_0^x g(s) ds$ and $k_0 = 1$. ■

Moreover, using an approach of [9] we obtain

Theorem 9 *Let us assume $f, g \in C^1(\mathbb{R}, \mathbb{R})$ with f bounded odd, and with g odd satisfying $G(u) \geq c(|u|^p - 1) \forall u \in \mathbb{R}$ and some constants $p > 2$, $c > 0$. Moreover we suppose*

$$\text{either } \cos \theta / T \notin \mathbb{Z} \text{ or } \sin \theta / T \notin \mathbb{Z} \quad (6.4)$$

along with the condition

$$\min_{k \in \mathbb{N}} \frac{4\pi^2}{T^2} k^2 \left(\nu^2 - g'(0) g_\theta \left(\frac{\pi k}{T} \right) \right) > f'(0). \quad (6.5)$$

Then equation (6.2) possesses a nonzero odd T -periodic solution.

Proof. We consider $\tilde{H}(x)$ over Y_1 . The condition (6.5) implies $\ker \text{Hess } \tilde{H}(0) = 0$ and $\text{index Hess } \tilde{H}(0) = 0$, so $0 \in Y_1$ is a strict local minimum of $\tilde{H}$. Next we take $x_0(z) = \sin \left(\frac{2\pi}{T} z \right)$. Then for any $\alpha > 0$ we have

$$\tilde{H}(\alpha x_0) \leq 2\nu^2 \alpha^2 \frac{\pi^2}{T} - \alpha^p \gamma_1 + \gamma_2 \alpha + \gamma_3 \quad (6.6)$$

for

$$\begin{aligned} \gamma_1 &:= c \int_0^T \left(\left| \sin \left(\frac{2\pi}{T} (z + \cos \theta) \right) - \sin \left(\frac{2\pi}{T} z \right) \right|^p + \left| \sin \left(\frac{2\pi}{T} (z + \sin \theta) \right) - \sin \left(\frac{2\pi}{T} z \right) \right|^p \right) dz \\ \gamma_2 &:= T \min_{\mathbb{R}} |f(x)|, \quad \gamma_3 := 2cT. \end{aligned}$$

Note $\gamma_1 > 0$ according to (6.4). Since $\tilde{H}(\alpha x_0) > 0$ for small $\alpha > 0$, (6.6) implies $\exists \alpha_0 > 0$, $\tilde{H}(\alpha_0 x_0) = 0$. Consequently, the mountain pass theorem [9, p. 141] can be applied for $\tilde{H}(x)$ to get its nonzero solution on Y_1 . The proof is finished. ■

For instance, if [32]

$$g(u) = \alpha_1 u + \alpha_2 u^3, \quad \alpha_{1,2} > 0, \quad f(u) = \omega \sin u, \quad (6.7)$$

where α_1, α_2 are the linear and nonlinear coupling coefficients in the longitudinal and transverse directions, respectively. Then, (6.4) and (6.5) are satisfied when $\tan \theta$ is irrational and $\frac{4\pi^2}{T^2} (\nu^2 - \alpha_1) \geq \omega$. So when $\omega \leq 0$, (6.2) with (6.7) possesses a nonzero odd T -periodic solution for any $\nu > \sqrt{\alpha_1}$ and any T, θ satisfying (6.4). This results can be related to [29] and [36, p. 274, Example (c)].

2.

Concerning the minimal period of a T -periodic solution of (2.4) we derive the following

Proposition 2 *Let f be globally Lipschitz continuous on $\mathbb{R}$, i.e. $\exists L > 0, |f(x) - f(y)| \leq L|x - y| \forall x, y \in \mathbb{R}$. Then a period T of any nonconstant T -periodic solution of (2.4) satisfies*

$$T \geq \frac{2\pi\nu}{\sqrt{L+8}}. \quad (6.8)$$

Proof. Let $P : L^2(0, T) \rightarrow L^2(0, T)$ be an orthogonal projection given by

$$PU := U(z) - \frac{1}{T} \int_0^T U(s) ds.$$

Let us denote by $G_\theta(U) : L^2(0, T) \rightarrow L^2(0, T)$ the right-hand side of (2.4). We note that G_θ has a global Lipschitz constant $L + 8$. Now we split

$$U(z) = U_0(z) + c, \quad c \in \mathbb{R}, \quad U_0 = PU.$$

We recall that the standard scalar product on $L^2(0, T)$ is denoted by $\langle \cdot, \cdot \rangle$ with the corresponding norm $\| \cdot \|$. Then (2.4) implies

$$\begin{aligned} \nu^2 \frac{4\pi^2}{T^2} \|U_0\|^2 &\leq -\nu^2 \langle U'', U_0 \rangle = -\langle G_\theta(U_0 + c), U_0 \rangle \\ &= -\langle P(G_\theta(U_0 + c) - G_\theta(c)), U_0 \rangle \leq (L + 8) \|U_0\|^2. \end{aligned} \quad (6.9)$$

Since U is nonconstant, we get $U_0 \neq 0$. Hence (6.9) gives (6.8). ■

3.

We are able to determine numerical profiles of found travelling waves only in results where constructive methods are used based on the Banach fixed point theorem, i.e. in Theorems 1, 7 and Corollary 1. We focus here on Theorem 1 assuming $\nu > 1$ and $LT^2 < 4\pi^2(\nu^2 - 1)$. Then there is a unique $x \in X_1$ of (3.3). Now like in [34, p. 270], for any $n \in \mathbb{N}$, we set $X_n :=$

$\text{span}\{e_k\}_{k=1}^n$. Let $P_n : X_1 \rightarrow X_1$ be the orthogonal projection onto X_n . Since $L_1 P_n = P_n L_1$, we consider an equation

$$L_1 x_n = P_n N_1(x_n), \quad x_n \in X_n. \quad (6.10)$$

Clearly $P N_1 : X_n \rightarrow X_n$ has a global Lipschitz constant L , and so $L_1^{-1} P N_1 : X_n \rightarrow X_n$ is contractive. Hence (6.10) has a unique solution $x_n \in X_n$. From (3.3) and (6.10) we derive

$$L_1(x - x_n) = P_n(N_1(x) - N_1(x_n)) + (I - P_n)N_1(x),$$

and so

$$\|x - x_n\| \leq L \|L_1^{-1}\| \|x - x_n\| + \|L_1^{-1}(I - P_n)N_1(x)\|.$$

That implies

$$\|x - x_n\| \leq \frac{\|L_1^{-1}(I - P_n)N_1(x)\|}{1 - \|L_1^{-1}\|L} \leq \frac{4\pi^2(\nu^2 - 1)}{4\pi^2(\nu^2 - 1) - T^2 L} \|L_1^{-1}(I - P_n)N_1(x)\|.$$

Note $\|L_1^{-1}(I - P_n)N_1(x)\| \rightarrow \infty$ as $n \rightarrow \infty$ and the convergence rapidity depends on a smoothness of the function $z \rightarrow N_1(x)(z) = f\left(\frac{2\pi}{T}z + x(z)\right)$.

For $f(x) = \sin x$, $T = 2\pi$, $\nu = 2$, $\theta = \pi/4$ and $x_n(z) = \frac{1}{\sqrt{2\pi}} \sum_{k=1}^n c_k \sin kz$, equation (6.10) has the form

$$c_k = \frac{1}{\sqrt{2\pi} \left(4k^2 - 8 \sin^2 \frac{\sqrt{2}}{4} k\right)} \int_0^{2\pi} \sin \left(z + \frac{1}{\sqrt{2\pi}} \sum_{k=1}^n c_k \sin kz \right) \sin kz \, dz, \quad k = 1, \dots, n. \quad (6.11)$$

We know that the right-hand side of (6.11) is a contraction, so it has a unique solution. Numerical profiles for $n = 15$ are drawn on Figure 3.

4.

For $\nu = 0$, equation (2.4) becomes a functional-difference equation for the ground-state of the FK model of the form

$$x(z + \cos \theta) + x(z - \cos \theta) + x(z + \sin \theta) + x(z - \sin \theta) - 4x(z) = f\left(\frac{2\pi}{T}z + x(z)\right). \quad (6.12)$$

(6.12) has complex behavior because it involves small divisors. This problem was rigorously solved in [4, 5], where it was proven that the hull function is monotonous increasing but it is not necessarily smooth and continuous. It is smooth only when the period of the ratio $2\pi/T$ is a "good" irrational number (according to Kolmogorov-Arnold-Moser theory) and when the potential $f(x)$ is not too large (incommensurate structures). For larger potentials or for commensurate structures, the hull function becomes purely discrete. These theorems were established for the 1d FK model but extend with practically the same proof in 2d and more




Figure 3: Numerical profiles of a uniform sliding state (3.1) of (2.4) (the monotonone one) and the corresponding odd T -periodic solution of (3.2), respectively, for $f(x) = \sin x$, $T = 2\pi$, $\nu = 2$, $\theta = \pi/4$ and $n = 15$ in (6.11).

dimensions in [6], thus including the 2d model treated in this paper. Thus a natural question arises: How the uniform sliding states which have continuous hull functions interpolate with the solutions at zero velocity were the hull functions may be either continuous or discontinuous? That is what is a relation between (3.2) and (6.12) as $\nu \rightarrow 0$? We can not answer in general on this question in this paper. On the other hand, singular delay differential equations are studied also in [22], and we believe that similar approach could be used to periodic travelling waves as well. More precisely, we consider (2.5) and we look for a $(1 + \xi\nu)$ -antiperiodic solution of it, i.e. satisfying (4.7) with $T = 1 + \xi\nu$, where $\xi \in \mathbb{R}$ is specified latter and ν is small. Then

$$w(z) := U(-\xi\nu z)$$

satisfies

$$\ddot{w}(z) + \xi^2 (w(z+1) + 2w(z) + w(z-1) + f(w(z))) = 0. \quad (6.13)$$

For the period of a τ -antiperiodic solution of (6.13), i.e. $w(z+\tau) = -w(z)$, we get the relation

$$\nu\xi(\tau - 1) = 1. \quad (6.14)$$

Then like in [22], we could apply the center manifold reduction using an approach of [15, 24, 25] near $w(z) \sim 0$ to derive a reduced ordinary differential equation. The next task would be to find a τ -antiperiodic solution of the reduced ordinary differential equation satisfying (6.14). We do not proceed with further computations, but we expect that square waves could be shown for (2.5) after embedding $f(u)$ into a 1-parametric family $f(u, \lambda)$, $\lambda \in \mathbb{R}$, $f(u, 0) = f(u)$, with certain conditions. The parameter ξ could be used to determine the dimension of the reduced o.d.e. in a spectral analysis similar to (A.1) in Appendix. So finally, we could show *square travelling waves* for (2.1) as $\nu \rightarrow 0$.

5.

We finish this paper with application of our method to a topological discrete sine-Gordon system [23] (an 1d lattice) given by

$$\ddot{\psi}_n = \alpha \cos \psi_n (\sin \psi_{n+1} + \sin \psi_{n-1}) - \left(\alpha + \frac{1}{2} \right) \sin \psi_n (\cos \psi_{n+1} + \cos \psi_{n-1}) \quad (6.15)$$

where $\alpha \in \mathbb{R}$ is a parameter. Inserting $\psi_n(t) = U(n - \nu t)$ into (6.15) we get

$$\nu^2 U''(z) = \alpha \cos U(z) (\sin U(z+1) + \sin U(z-1)) - \left(\alpha + \frac{1}{2} \right) \sin U(z) (\cos U(z+1) + \cos U(z-1)). \quad (6.16)$$

Following the proof of Theorem 5 we have,

Theorem 10 *For any $\nu > 0$, $T > 0$ and $\alpha \in \mathbb{R}$ satisfying*

$$\frac{4\pi^2\nu^2}{T^2} - 4\alpha \sin^2 \frac{\pi}{T} < 1, \quad (6.17)$$

(6.16) possesses a nonzero odd T -periodic solution.

Proof. Now we take the function

$$\check{H}(x) := \int_0^T \left\{ \nu^2 \frac{x'(z)^2}{2} + \alpha \sin U(z+1) \sin U(z) + \left(\alpha + \frac{1}{2} \right) \cos U(z) \cos U(z-1) \right\} dz$$

and $k_0 = 1$ in the proof of Theorem 5. ■

We note [23] that the "continuum limit" of (6.16) is $\alpha \rightarrow -\infty$. Then condition (6.17) holds for $\alpha < 0$ and $T > 2\pi\sqrt{\nu^2 - \alpha}$. On the other hand, the "anti-continuum limit" of (6.16) is $\alpha \rightarrow 0$. Then condition (6.17) holds for $|\alpha| < 1/4$ and

$$T > \frac{2\pi\nu}{\sqrt{1 - 4|\alpha|}}$$

with $\nu > 0$.

Acknowledgments

This work was supported by the IACM FORTH Greece and the Grant VEGA-SAV 2/4135/25. We also thanks to the referees for valuable comments and suggestions which substantially improved our paper.

A Spectral analysis of certain linear operators

The spectrums of the above linear self-adjoint bounded operators can be easily computed, since standard orthogonal basis of the corresponding functional Hilbert spaces form their eigenvectors. This follows from the next facts:

1. - these orthogonal basis are created by functions of a form $c_1 \cos a_1 z + c_2 \sin a_1 z$ for constants $c_{1,2}, a_1$,
2. - these linear operators are linear combinations of the shift mapping $x(z) \rightarrow x(z + a_2)$ and the differential operator $x(z) \rightarrow x''(z)$ for a constant a_2 ,
3. - subspaces $\text{span} \{ \cos a_1 z, \sin a_1 z \}$ are invariant under operators from point 2.

For instance, in Theorem 1, the standard basis $\{e_k\}$ of X_1 are readily verified to be also eigenvectors of L_1 .

On the other hand, we have also another spectrum associated to (3.2). Following [15, 24, 25], we can write (3.2) as an evolution equation

$$\begin{aligned} \dot{x}(t) &= \xi(t) \\ \dot{\xi}(t) &= -\frac{4}{\nu^2}x(t) + \frac{1}{\nu^2} \left(X(t, \cos \theta) + X(t, -\cos \theta) \right. \\ &\quad \left. + X(t, \sin \theta) + X(t, -\sin \theta) \right) - \frac{1}{\nu^2} f\left(\frac{2\pi}{T}t + x(t)\right) \\ X_t(t, v) &= X_v(t, v) \end{aligned} \tag{A.1}$$

on the space

$$H := \left\{ (x, \xi, X) \in \mathbb{R}^2 \times C^1([-1, 1], \mathbb{R}) \mid X(0) = x \right\}.$$

The linear part of (A.1) is

$$L = \begin{pmatrix} 0, & 1, & 0 \\ -\frac{4}{\nu^2}, 0, \frac{1}{\nu^2} \left(\partial^{\cos \theta} + \partial^{-\cos \theta} + \partial^{\sin \theta} + \partial^{-\sin \theta} \right) \\ 0, & 0, & \partial_v \end{pmatrix}.$$

Here $\partial^a X = X(a)$ and ∂_v is the differentiation with respect to v . The spectrum of L is given by the equation

$$\lambda^2 + \frac{1}{\nu^2} \left(4 - 2 \cosh(\lambda \cos \theta) - 2 \cosh(\lambda \sin \theta) \right) = 0. \tag{A.2}$$

The imaginary eigenvalues $\lambda = \imath q$ are determined by

$$\nu^2 q^2 - 4 \sin^2 \left(\frac{q \cos \theta}{2} \right) - 4 \sin^2 \left(\frac{q \sin \theta}{2} \right) = 0. \tag{A.3}$$

Since the nonlinear part of (A.1) is T -periodic, we see that condition i) of Theorem 1 are just nonresonance conditions between the linear and forcing/nonlinear parts of (A.1). This should give an alternative proof of Theorem 1.

References

- [1] AIGNER, A. A., CHAMPNEYS, A. R. AND ROTHOS, V. M.: *A new barrier to the existence for moving kinks in Frenkel-Kontorova lattices*, Physica D **186** (3-4) (2003), 148–170.
- [2] ARIOLI, G. AND GAZZOLA, F.: *Periodic motions of an infinite lattice of particles with nearest neighbor interaction*, Nonl. Anal. **26** (1996), 1103–1114.
- [3] AUBRY, S.: *Breathers in nonlinear lattices: existence, linear stability and quantization*, Physica D **103** (1997), 201–250.
- [4] AUBRY, S.: *The twist map, the extended Frenkel-Kontorova model and the devil's staircase*, Physica D **7** (1983), 240–258.
- [5] AUBRY, S. AND LE DAERON, P. Y.: *The discrete Frenkel-Kontorova model and its extensions*, Physica D **8** (1983), 381–422.
- [6] AUBRY, S., GOSSO, J. P., ABRAMOVICI, G., RAIMBAULT, J. L. AND QUEMERAIS, P.: *Effective discommensurations in the incommensurate ground states of extended Frenkel-Kontorova models*, Physica D **47** (1991), 461–497.
- [7] BROWN, R. F.: *A Topological Introduction to Nonlinear Analysis*, Birkhäuser, Boston, 1993.
- [8] CESARI, L.: *Asymptotic behavior and stability problems in ordinary differential equations*, Springer-Verlag, Berlin, 1959.
- [9] CHOW, S. N. AND HALE, J. K.: *Methods of Bifurcation Theory*, Springer-Verlag, New York, 1982.
- [10] CHOW, S. N., MALLET-PARET, J. AND SHEN, W.: *Traveling waves in lattice dynamical systems*, J. Differential Equations **149** (1998), 248–291.
- [11] COULLET, P. AND WALGRAEF, D.: *Spatial forcing of 2d wave patterns*, Europhys. Lett. **10** (1989), 525–531.
- [12] DMITRIEV, S. V., KEVREKIDIS, P. G., YOSHIKAWA, N. AND FRANTZESKAKIS, D. J.: *Exact static solutions for discrete ϕ^4 models free of the Peierls-Nabarro barrier: discretized first integral approach*, preprint, arXiv:nlin:PS/0603074 v2 7 Aug 2006.
- [13] FALO, F., BISHOP, A. R., LOMDAHL, P. S. AND HOROVITZ, B.: *Langevin molecular dynamics of interfaces: Nucleation versus spiral growth*, Phys. Rev. B **43**(10) (1991), 8081–8088.
- [14] FEČKAN, M.: *Nontrivial critical points of asymptotically quadratic functions at resonance*, Ann. Polonici Math. **67** (1997), 43–57.

- [15] FEČKAN, M. AND ROTHOS, V. M.: *Kink-like periodic travelling waves for lattice equations with on-site and intersite potentials*, Dynamics of PDEs **2**(4) (2005), 357–370.
- [16] FLACH, S. AND WILLIS, C. R.: *Discrete breather*, Phys. Rep. **295** (1998), 181–264.
- [17] FLORÍA, L. M. AND FALO, F.: *Shapiro steps in the steady-state dynamycs of imcommensurate structures*, Phys. Rev. Letters **68**(18) (1992), 2713–2718.
- [18] FRIESECKE, G. AND MATTHIES, K.: *Geometric solitay waves in a 2d mass-spring lattice*, Disc. Cont. Dyn. Syst.-B **3** (2003), 105–114.
- [19] FRIESECKE, G. AND PEGO, R. L.: *Solitay waves on FPU lattices: I. Qualitative properties, renolmalization and continuun limit*, Nonlinearity **12** (1999), 1601–1627.
- [20] FRIESECKE, G. AND PEGO, R. L.: *Solitay waves on FPU lattices: II. Linear implies nonlinear stability*, Nonlinearity **15** (2002), 1343–1359.
- [21] FRIESECKE, G. AND WATTIS, J.: *Existence theorem for solitary waves on lattices*, Comm. Math. Phys. **161** (1994), 391–418.
- [22] HALE, J. K. AND HUANG, W. M.: *Period doubling in singularly perturbed delay equations*, J. Differential Equations **114** (1994), 1–23.
- [23] HASKINS, M. AND SPEIGHT, J. M.: *Breathers in the weakly coupled topological discrete sine-Gordon system*, Nonlinearity **11** (1998), 1651–1671.
- [24] IOOSS, G.: *Trawelling waves in the Fermi-Pasta Ulam lattice*, Nonlinearity **13**(3) (2000), 849–866.
- [25] IOOSS, G. AND KIRCHGÄSSNER, K.: *Traveling waves in a chain of coupled nonlinear oscillators*, Comm. Math. Phys. **211**(2) (2000), 439–464.
- [26] KARTIEL, G.: *Existence of travelling waves in discrete sine-Gordon rings*, SIAM J. Math. Anal. **36** (2005), 1433–1443.
- [27] MAWHIN, J.: *Contractive mappings and periodically perturbed conservative systems*, Arch. Math. **12** (1976), 67–74.
- [28] MAWHIN, J. AND WILLEM, M.: *Critical Point Theory and Hamiltonian Systems*, Springer, New York, 1989.
- [29] PANKOV, A. AND PFLÜGER, K.: *Traveling waves in lattice dynamical systems*, Math. Meth. Appl. Sci. **23** (2000), 1223–1235.
- [30] PEYRARD M. AND KRUSKAL, M. D.: *Kink dynamics in the highly discrete sine-Gordon system*, Physica D **14** (1984), 88–102.

- [31] PEYRARD M., PNEVMATIKOS, ST. AND FLYTANIS, N.: *Dynamics of two-component solitary waves in hydrogen-bonded chains*, Phys. Rev. A **36** (1987), 903–914.
- [32] POUGET, J., REMOISSENET, M. AND TAMGA, J. M.: *Energy self-localization and gap local pulses in a two-dimensional nonlinear lattice*, Phys. Rev. B **47**(22) (1993), 14 866–14874.
- [33] RUDIN, W.: *Real and Complex Analysis*, McGraw-Hill, Inc., New York, 1974.
- [34] SAVIN, A. V, ZOLOTARYUK, Y. AND EILBECK, J. C.: *Moving kinks and nonopterons in the nonlinear Klein-Gordon lattice*, Physica D **138** (2000), 267–281.
- [35] SCOTT, A.: *Nonlinear Science: Emergence and Dynamics of Coherent Structures*, Oxford Univ. Press, Oxford, 2003.
- [36] SMETS, D. AND WILLEM, M.: *Solitary waves with prescribed speed on infinite lattices*, J. Funct. Anal. **149** (1997), 266–275.
- [37] SNEDDON, L.: *Sliding dynamics of the incommensurate chain*, Phys. Rev. Letters **52**(1) (1984), 65–68.
- [38] STRUNZ, T. AND ELMER, F. J.: *Driven Frenkel-Kontorova model. I. Uniform sliding states and dynamical domains of different particle densities*, Phys. Review E **58**(2) (1998), 1601–1611.
- [39] TAMGA, J. M, REMOISSENET, M. AND POUGET, J.: *Breathing solitary waves in a sine-Gordon two dimensional lattice*, Phys. Rev. Letters **75**(3) (1995), 357.
- [40] TITCHMARSH, E. C.: *The Theory of Functions*, 2nd ed., Oxford Univ. Press, Fair Lawn, N. J., 1939.